arXiv:1010.5535v1 [math.AP] 26 Oct 2010

A STRENGTHENING OF THE ENERGY INEQUALITY FOR THE LERAY-HOPF SOLUTIONS OF THE 3D PERIODIC NAVIER-STOKES EQUATIONS

R. DASCALIUC

Abstract. In present note we establish the following inequality for the the Leray-Hopf solutions of the 3-D Ω-periodic Navier-Stokes Equations:

$$\phi(|u(t)|^2)-\phi(|u(t_0)|^2)\le 2\int_{t_0}^{t}\phi'(|u(\tau)|^2)\left[-\nu|A^{1/2}u(\tau)|^2+(g(\tau),u(\tau))\right]d\tau$$

for all t_0 Leray-Hopf points, $t\ge t_0$, and $\phi:\mathbb{R}_+\to\mathbb{R}$ is an absolutely continouos non-decreasing function with bounded derivative. Here $(\cdot,\cdot)$ and $|\cdot|$ is correspondingly the L^2 inner product and the L^2 norm on Ω, and A is the Stokes operator.

1. PRELIMINARIES

We consider three dimensional incompressible Navier-Stokes equations (NSE)

$$\begin{aligned}\frac{\partial}{\partial t}\mathbf{u}(t,\mathbf{x})-\nu\Delta\mathbf{u}(t,\mathbf{x})+(\mathbf{u}(t,\mathbf{x})\cdot\nabla)\mathbf{u}(t,\mathbf{x})+\nabla p(t,\mathbf{x})&=\mathbf{f}(t,\mathbf{x})\\ \nabla\cdot\mathbf{u}(t,\mathbf{x})&=0,\end{aligned}\tag{1.1}$$

where $\mathbf{x}\in\mathbb{R}^3$, $t\in\mathbb{R}$; and $\mathbf{u}(t,\mathbf{x}),\mathbf{f}(t,\mathbf{x})\in\mathbb{R}^3$ and $p(t,\mathbf{x})\in\mathbb{R}$ for all t and $\mathbf{x}$. We supplement (1.1) with periodic boundary conditions

$$\begin{gathered}\mathbf{u},p,\mathbf{f}\text{ are }\Omega-\text{periodic},\\ \int_\Omega\mathbf{u}(t,\mathbf{x})\,d\mathbf{x}=0\text{ and }\int_\Omega\mathbf{f}(t,\mathbf{x})\,d\mathbf{x}=0\text{ for all }t,\end{gathered}\tag{1.2}$$

where

$$\Omega=[0,L]^3,$$

and the initial condition

$$\begin{gathered}\mathbf{u}(0,\mathbf{x})=\mathbf{u}_0(\mathbf{x})\text{ for all }\mathbf{x},\\ \mathbf{u}_0\text{ satisfies (1.2).}\end{gathered}\tag{1.3}$$

Taking the Leray projector P_L of (1.1) we obtain

$$\frac{d}{dt}\mathbf{u}+\nu A\mathbf{u}+B(\mathbf{u},\mathbf{u})=\mathbf{g},\tag{1.4}$$

where

$$A\mathbf{u}=-P_L\Delta\mathbf{u},\ B(\mathbf{u},\mathbf{u})=P_L(\mathbf{u}\cdot\nabla)\mathbf{u},\ \text{and }\mathbf{g}=P_L\mathbf{f}.$$

We introduce the following functional space

$$H=\left\{\mathbf{u}:\mathbf{u}\ -\ \Omega-\text{periodic},\ \mathbf{u}\in L^2(\Omega)^3,\ \nabla\cdot\mathbf{u}=0,\int_\Omega\mathbf{u}=0\right\}\tag{1.5}$$

Date: October 28, 2010.

For any $\mathbf{u}, \mathbf{v} \in H$ denote

$$(\mathbf{u}, \mathbf{v}) = \int_\Omega \mathbf{u} \cdot \mathbf{v}$$

and

$$|\mathbf{u}|^2 = \int_\Omega \mathbf{u} \cdot \mathbf{u},$$

the L^2 inner product and norm on H. Note that $A : D(A) \in H \to H$ is a positive self adjoint operator with a compact inverse; its domain $D(A)$ is dense in H. Denote by λ_k, $k \in \mathbb{N}$, its eigenvalues arranged in the increasing order and counting the multiplicities. We also introduce the functional space

$$V = \left\{\mathbf{u} \in H : \ \mathbf{u} \in H^1(\Omega)^3\right\} = \left\{\mathbf{u} \in H : \ \mathbf{u} \in D(A^{1/2})\right\}. \tag{1.6}$$

Define the inner product and the norm on V by

$$((\mathbf{u}, \mathbf{v})) = (A^{1/2}\mathbf{u}, A^{1/2}\mathbf{v})$$

and

$$||\mathbf{u}||^2 = |A^{1/2}\mathbf{u}|^2$$

for all $\mathbf{u}, \mathbf{v} \in V$. Note that

$$\lambda_1 |\mathbf{u}|^2 \leq ||\mathbf{u}||^2$$

for all $\mathbf{u} \in V$.

The bilinear operator B has the following orthogonality property:

$$(B(\mathbf{u}, \mathbf{v}), \mathbf{w}) = -(B(\mathbf{u}, \mathbf{w}), \mathbf{v}), \text{ for all } \mathbf{u}, \mathbf{v}, \mathbf{w} \in V,$$

and consequently

$$(B(\mathbf{u}, \mathbf{v}), \mathbf{v}) = 0, \text{ for all } \mathbf{u}, \mathbf{v} \in V.$$

In this paper we study weak solutions of the NSE, i.e. H-valued functions $\mathbf{u}(t)$ that satisfy

$$\frac{d}{dt}(\mathbf{u}(t), \mathbf{v}) + \nu((\mathbf{u}(t), \mathbf{v})) + (B(\mathbf{u}(t), \mathbf{u}(t)), \mathbf{v}) = (\mathbf{g}(t), \mathbf{v}) \text{ for all } \mathbf{v} \in V.$$

For more background on the Leray-Hopf solutions of Navier-Stokes equations and on the functional setting used here consult [1, 3, 2, 7, 6, 5, 4]

The principal result about the existence of the weak (Leray-Hopf) solutions of the NSE can be stated as follows.

Theorem 1.1. *Let $\mathbf{u}_0 \in H$ and $\mathbf{g} \in L^2([0,T], H)$, where $T > 0$ given. Then there exist a weak solution $\mathbf{u} \in L^2([0,T], V)$, which is weakly continuous as a function from $[0,T]$ to H.*

Moreover, the following inequality holds

$$|\mathbf{u}(t)|^2 - |\mathbf{u}(t_0)|^2 \leq 2 \int_{t_0}^{t} \left[-\nu ||\mathbf{u}(\tau)||^2 + (\mathbf{g}(\tau), \mathbf{u}(\tau))\right] \, d\tau \tag{1.7}$$

for all $t_0 \geq 0$ Lebesgue points of $|\mathbf{u}(\tau)|^2$ and all $t \in [t_0, T]$.

In fact, the set of Lebesgue points of $|\mathbf{u}(\tau)|^2$ is dense in $[0,T]$ and includes $t_0 = 0$ and any t_0 such that $\mathbf{u}(t_0) \in V$.

Under the additional assumptions on the initial data, one gets local existence of the regular solutions of the NSE.

Theorem 1.2. *Suppose* $\mathbf{u}_0 \in V$ *and* $\mathbf{g} \in L^2([0,T],H)$. *Then there exists* $T_* = T_*(||u_0||,\nu,\Omega) > 0$ *such that the weak solution* $\mathbf{u}$ *from Theorem 1.1 is unique on* $[0,T_*]$ *and* $\mathbf{u} \in L^2([0,T_*],D(A)) \cap C([0,T_*],V)$ *(i.e.* $\mathbf{u}$ *is* regular *on* $[0,T_*]$*). Moreover, the inequality* (1.7) *becomes equality for all* $t_0, t \in [0,T_*]$, $t \geq t_0$.

It is important to mention that any weak solution of the NSE is regular on the set

$$\mathcal{G} = \cup_n I_n,$$

where $\{I_n\}_n$ is a countable family of disjoint intervals with $\mathrm{cl}(\mathcal{G}) = [0,T]$, and the fractal dimension of $[0,T]\backslash\mathcal{G}$ less or equal to 1/2.

2. Proof of the main result

Theorem 2.1. *Let* $\mathbf{g} \in L^2([0,T],H)$ *and suppose* $\mathbf{u}$ *is a weak solution for the NSE. Let* $\phi : \mathbb{R}_+ \to \mathbb{R}$ *be an absolutely continouos non-decreasing function with* $\phi' \in L^\infty(\mathbb{R}_+,\mathbb{R}_+)$. *Then the following inequality holds*

$$\phi(|\mathbf{u}(t)|^2) - \phi(|\mathbf{u}(t_0)|^2) \leq 2\int_{t_0}^{t} \phi'(|\mathbf{u}(\tau)|^2)\left[-\nu||\mathbf{u}(\tau)||^2 + (\mathbf{g}(\tau),\mathbf{u}(\tau))\right]\,d\tau \tag{2.1}$$

for all $t_0 \in [0,T]$ *- Lebesgue point for* $|\mathbf{u}(\tau)|^2$ *and all* $t \in [0,T]$, $t \geq t_0$.

Proof. Let $t_0, t \in [0,T]$, $t \geq t_0$, t_0 - Lebesgue point for $|\mathbf{u}(\tau)|^2$. Let

$$M = \mathrm{essup}_{\tau\in[t_0,t]}\phi'(|\mathbf{u}(\tau)|^2)$$

Choose an $\epsilon > 0$. Then there exists $\delta = \delta(\epsilon) > 0$ such that, for any countable family of disjoint intervals $\{[t_n,\tau_n]\} \subset [t_0,t]$ such that $\sum_n(\tau_n - t_n) < \delta$,

$$M \int\limits_{\cup_n[t_n,\tau_n]} \frac{|\mathbf{g}(\tau)|^2}{\nu\lambda_1} d\tau < \epsilon. \tag{2.2}$$

and

$$\left| \int\limits_{\cup_n[t_n,\tau_n]} \phi'(|\mathbf{u}(\tau)|^2)\left[-\nu||\mathbf{u}(\tau)||^2 + (\mathbf{g}(\tau),\mathbf{u}(\tau))\right]\,d\tau \right| < \epsilon \tag{2.3}$$

Recall that $\mathbf{u}$ is regular on a set

$$\mathcal{G} = \cup_n I_n,$$

where $\mathrm{int}(I_n) = (\alpha_n,\beta_n)$ and $\sum_n(\beta_n - \alpha_n) = t - t_0$. Clearly, inside each of the intervals (α_n,β_n) we have

$$\phi(|\mathbf{u}(s_1)|^2) - \phi(|\mathbf{u}(s_0)|^2) = 2\int_{s_0}^{s_1} \phi'(|\mathbf{u}(\tau)|^2)\left[-\nu||\mathbf{u}(\tau)||^2 + (\mathbf{g}(\tau),\mathbf{u}(\tau))\right]\,d\tau$$

for all $s_0, s_1 \in (\alpha_n,\beta_n)$, $s_0 \leq s_1$.

Note that there exists $N = N(\delta)$ such that

$$\sum_{n=1}^{N}(\beta_n - \alpha_n) \geq t - t_0 - \delta/2.$$

Re-arrange these first N intervals such that $\beta_n \leq \alpha_{n+1}$ for all $n = 1..N-1$. Choose $\tau_{n-1}, t_n \in (\alpha_n, \beta_n)$, $n = 1..N$ and $\tau_N = t$, so that $\tau_{n-1} < t_n < \tau_n$ for $n = 1..N$ and

$$\sum_{n=0}^{N} (\tau_n - t_n) < \delta.$$

Then, by the (1.7),

$$|\mathbf{u}(\tau_n)|^2 - |\mathbf{u}(t_n)|^2 \leq 2 \int_{t_n}^{\tau_n} \left[-\nu ||\mathbf{u}(\tau)||^2 + (\mathbf{g}(\tau), \mathbf{u}(\tau))\right] d\tau,$$

for all $n = 0..N$. Then,

$$|\mathbf{u}(\tau_n)|^2 - |\mathbf{u}(t_n)|^2 \leq |\mathbf{u}(\tau_n)|^2 - |\mathbf{u}(t_n)|^2 + \nu \int_{t_n}^{\tau_n} ||\mathbf{u}(\tau)||^2 d\tau \leq \int_{t_n}^{\tau_n} \frac{|\mathbf{g}(\tau)|^2}{\nu\lambda_1} d\tau, \tag{2.4}$$

for all $n = 0..N$. On the other hand, since ϕ is increasing,

$$\phi(|\mathbf{u}(\tau_n)|^2) - \phi(|\mathbf{u}(t_n)|^2) \leq \begin{cases} 0, \text{ if } |\mathbf{u}(\tau_n)| < |\mathbf{u}(t_n)|; \\ M(|\mathbf{u}(\tau_n)|^2 - |\mathbf{u}(t_n)|^2), \text{ otherwise.} \end{cases}$$

Use (2.4) to obtain

$$\phi(|\mathbf{u}(\tau_n)|^2) - \phi(|\mathbf{u}(t_n)|^2) \leq M \int_{t_n}^{\tau_n} \frac{|\mathbf{g}(\tau)|^2}{\nu\lambda_1} d\tau.$$

Thus, taking into the account (2.2) and (2.3), we can estimate

$$\begin{aligned} &\phi(|\mathbf{u}(t)|^2) - \phi(|\mathbf{u}(t_0)|^2) \\ &= \sum_{n=0}^{N} \left(\phi(|\mathbf{u}(\tau_n)|^2) - \phi(|\mathbf{u}(t_n)|^2)\right) + \sum_{n=1}^{N} \left(\phi(|\mathbf{u}(t_n)|^2) - \phi(|\mathbf{u}(\tau_{n-1})|^2)\right) \\ &\leq M \int_{\cup_n [t_n, \tau_n]} \frac{|\mathbf{g}(\tau)|^2}{\nu\lambda_1} d\tau + 2 \sum_{n=1}^{N} \int_{\tau_{n-1}}^{t_n} \Psi(\tau) d\tau \\ &\leq \epsilon + \sum_{n=1}^{N} \int_{\tau_{n-1}}^{t_n} \Psi(\tau) d\tau \leq 2\epsilon + 2 \int_{t_0}^{t} \Psi(\tau) d\tau \end{aligned} \tag{2.5}$$

where

$$\Psi(\tau) = \phi'(|\mathbf{u}(\tau)|^2) \left[-\nu ||\mathbf{u}(\tau)||^2 + (\mathbf{g}(\tau), \mathbf{u}(\tau))\right].$$

Returning to (2.5), make $\epsilon \to 0$ to obtain (2.1).

□

Corollary 2.1. *Under the assumptions of Theorem 2.1, let* $\psi \in C^1(\mathbb{R}_+, \mathbb{R})$ *be such that* $\psi'(\xi) \leq 0$ *for all* $\xi \geq 0$. *Then the following inequality holds*

$$\psi(|\mathbf{u}(t)|^2) - \psi(|\mathbf{u}(t_0)|^2) \geq 2 \int_{t_0}^{t} \psi'(|\mathbf{u}(\tau)|^2) \left[-\nu ||\mathbf{u}(\tau)||^2 + (\mathbf{g}(\tau), \mathbf{u}(\tau))\right] d\tau \tag{2.6}$$

for all $t_0 \in [0, T]$ *- Lebesgue point for* $|\mathbf{u}(\tau)|^2$ *and all* $t \in [0, T]$, $t \geq t_0$.

Proof. This result follows by applying Theorem 2.1 to $\phi = -\psi$. □

ACKNOWLEDGEMENTS

The author would like to thank Professor Ciprian Foias for helpful suggestions and comments.

REFERENCES

[1] P. Constantin and C. Foias. *Navier-Stokes equations.* Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 1988.

[2] C. Doering and J. Gibbon. *Applied analysis of the Navier-Stokes equations.* Cambridge Texts in Applied Mathematics. Cambridge University Press, Cambridge, 1995.

[3] C. Foias, O. Manley, R. Rosa, and R. Temam. *Navier-Stokes equations and turbulence*, volume 83 of *Encyclopedia of Mathematics and its Applications.* Cambridge University Press, Cambridge, 2001.

[4] P. G. Lemarié-Rieusset. *Recent developments in the Navier-Stokes problem*, volume 431 of *Chapman & Hall/CRC Research Notes in Mathematics.* Chapman & Hall/CRC, Boca Raton, FL, 2002.

[5] Hermann Sohr. *The Navier-Stokes equations.* Birkhäuser Advanced Texts: Basler Lehrbücher. [Birkhäuser Advanced Texts: Basel Textbooks]. Birkhäuser Verlag, Basel, 2001. An elementary functional analytic approach.

[6] R. Temam. *Navier-Stokes equations and nonlinear functional analysis*, volume 66 of *CBMS-NSF Regional Conference Series in Applied Mathematics.* Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, second edition, 1995.

[7] R. Temam. *Navier-Stokes equations.* AMS Chelsea Publishing, Providence, RI, 2001. Theory and numerical analysis, Reprint of the 1984 edition.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF VIRGINIA, CHARLOTTESVILLE, VA 22904